

**Joint Replacement, Sports Medicine,  
Fracture Care & General Orthopaedics**

Robert F. Jones, MD  
Marisa A. Schmitt, MD  
Alison Bird, MHP, PA-C



**Phone (978) 534-6333  
www.longviewortho.com**

100 Hospital Road, Suite 3C  
Leominster, MA 01453  
Fax (978) 840-0966

## **HAND SURGERY - Postoperative Instructions**

- Ganglion**
- Trigger Finger**
- deQuervaine's Release**

**Follow-up appointment:** Our office will call you the next business day after surgery to schedule a postoperative visit (usually 10 days postop). At that time you will be given the opportunity to ask questions about your surgery or discuss any difficulties you may have.

**About your dressing:** Your hand is covered with a sterile dressing following the procedure, and this should be kept clean and dry for 48 hours. You may shower or bathe, but be sure that the dressing is covered with a plastic bag and out of the flow of water. You may change the dressing at any time if it becomes soiled or wet. Remove the dressing after TWO days and cover the wound with gauze or a large band aid. The stitches used to close the wounds are made of nylon and will be removed when you return for your follow-up visit.

**About pain and swelling:** Some discomfort is to be expected following this surgery. Your surgeon uses a local anesthesia surrounding the incision that will provide pain relief for 6-10 hours following surgery. The best way to control the discomfort is to keep the hand elevated (above the level of the heart) and apply ice (enclosed in a plastic bag and wrapped in a dry towel) to the hand for 20 minutes out of an hour, repeating as needed.

For pain, use the narcotic medication that your doctor ordered. Please do not combine narcotics with alcoholic beverages.

Nonsteroidal anti-inflammatories (like Motrin, Advil or Aleve) may be used with or instead of the narcotic medication.

**Postoperative activities:** The fingers are left free so they can be moved as much as possible. Using the fingers immediately will speed up your recovery and help the pain resolve. Begin to move the wrist and hand more as the pain subsides. Gradually advance activities as tolerated.

**Return to driving/work:** Please do not drive for 24 hours following surgery. **OK to resume driving when you are no longer taking narcotics and can comfortably grip the steering wheel.** Return to work will depend on your work duties- this should be discussed with your surgeon.

**Warning signs:** Please call our office at (978)534-6333 for fevers, chills, excess drainage, severe pain or any questions.